



Bibliografía

Referencias

1. EFSA. Scientific Opinion on Q fever. EFSA J. (2010) 8:1595. doi: 10.2903/j.efsa.2010.1595
2. van den Brom R, van EE, Roest HI, van der Hoek W, Vellema P. Coxiella burnetii infections in sheep or goats: an opinionated review. Vet Microbiol. (2015) 181:119–29. doi: 10.1016/j.vetmic.2015.07.011
3. Agerholm JS. Coxiella burnetii associated reproductive disorders in domestic animals—a critical review. Acta Vet Scand. (2013) 55:13. doi: 10.1186/1751-0147-55-13
4. Alvarez-Alonso R, Basterretxea M, Barandika JF, Hurtado A, Idiazabal J, Jado I, et al. A Q fever outbreak with a high rate of abortions at a dairy goat farm: Coxiella burnetii shedding, environmental contamination, and viability. Appl Environ Microbiol. (2018) 84:e01650–18. doi: 10.1128/AEM.01650-18
5. Astobiza I, Barandika JF, Hurtado A, Juste RA, García-Pérez AL. Kinetics of Coxiella burnetii excretion in a commercial dairy sheep flock after treatment with oxytetracycline. Vet J. (2010) 184:172–5. doi: 10.1016/j.tvjl.2009.01.017
6. Berri M, Souriau A, Crosby M, Crochet D, Lechopier P, Rodolakis A. Relationships between the shedding of Coxiella burnetii, clinical signs and serological responses of 34 sheep. Vet Rec. (2001) 148:502–5. doi: 10.1136/vr.148.16.502
7. Berri M, Rousset E, Hechard C, Champion JL, Dufour P, Russo P, et al. Progression of Q fever and Coxiella burnetii shedding in milk after an outbreak of enzootic abortion in a goat herd. Vet Rec. (2005) 156:548–9. doi: 10.1136/vr.156.17.548
8. Berri M, Rousset E, Champion JL, Russo P, Rodolakis A. Goats may experience reproductive failures and shed Coxiella burnetii at two successive parturitions after a Q fever infection. Res Vet Sci. (2007) 83:47–52. doi: 10.1016/j.rvsc.2006.11.001
9. Joulé A, Laroucau K, Bailly X, Prigent M, Gasqui P, Lepetitcolin E, et al. Circulation of Coxiella burnetii in a naturally infected flock of dairy sheep: shedding dynamics, environmental contamination, and genotype diversity. Appl Environ Microbiol. (2015) 81:7253–60. doi: 10.1128/AEM.02180-15





10. Rodolakis A, Berri M, Hechard C, Caudron C, Souriau A, Bodier CC, et al. Comparison of *Coxiella burnetii* shedding in milk of dairy bovine, caprine, and ovine herds. *J Dairy Sci.* (2007) 90:5352–60. doi: 10.3168/jds.2006-815
11. Rousset E, Durand B, Champion JL, Prigent M, Dufour P, Forfait C, et al. Efficiency of a phase 1 vaccine for the reduction of vaginal *Coxiella burnetii* shedding in a clinically affected goat herd. *Clin Microbiol Infect.* (2009) 15(Suppl 2):188–9. doi: 10.1111/j.1469-0691.2008.02220.x
12. Dijkstra F, van der Hoek W, Wijers N, Schimmer B, Rietveld A, Wijkman CJ, et al. The 2007–2010 Q fever epidemic in The Netherlands: characteristics of notified acute Q fever patients and the association with dairy goat farming. *FEMS Immunol Med Microbiol.* (2012) 64:3–12. doi: 10.1111/j.1574-695X.2011.00876.x
13. Alvarez-Alonso R, Barandika JF, Ruiz-Fons F, Ortega-Araiztegi I, Jado I, Hurtado A, et al. Stable levels of *Coxiella burnetii* prevalence in dairy sheep flocks but changes in genotype distribution after a 10-year period in northern Spain. *Acta Vet Scand.* (2018) 60:75. doi: 10.1186/s13028-018-0429-x
14. Barlozzari G, Sala M, Iacoponi F, Volpi C, Polinori N, Rombola P, et al. Cross-sectional serosurvey of *Coxiella burnetii* in healthy cattle and sheep from extensive grazing system in central Italy. *Epidemiol Infect.* (2020) 148:e9. doi: 10.1017/S0950268819002115
15. Muema J, Thumbi SM, Obonyo M, Wanyoike S, Nanyingi M, Osoro E, et al. Seroprevalence and factors associated with *Coxiella burnetii* infection in small ruminants in baringo County, Kenya. *Zoonoses Public Health.* (2017) 64:e31–43. doi: 10.1111/zph.12342
16. Plummer PJ, McClure JT, Menzies P, Morley PS, van den Brom R, Van Metre DC. Management of *Coxiella burnetii* infection in livestock populations and the associated zoonotic risk: a consensus statement. *J Vet Intern Med.* (2018) 32:1481–94. doi: 10.1111/jvim.15229
17. Rizzo F, Vitale N, Ballardini M, Borromeo V, Luzzago C, Chiavacci L, et al. Q fever seroprevalence and risk factors in sheep and goats in northwest Italy. *Prev Vet Med.* (2016) 130:10–7. doi: 10.1016/j.prevetmed.2016.05.014
18. Astobiza I, Barandika JF, Ruiz-Fons F, Hurtado A, Povedano I, Juste RA, et al. *Coxiella burnetii* shedding and environmental contamination at lambing in two highly naturally-infected dairy sheep flocks after vaccination. *Res Vet Sci.* (2011) 91:e58–63. doi: 10.1016/j.rvsc.2010.11.014
19. Berri M, Souriau A, Crosby M, Rodolakis A. Shedding of *Coxiella burnetii* in ewes in two pregnancies following an episode of *Coxiella* abortion in a sheep flock. *Vet Microbiol.* (2002) 85:55–60. doi: 10.1016/S0378-1135(01)00480-1



20. Astobiza I, Barandika JF, Ruiz-Fons F, Hurtado A, Povedano I, Juste RA, et al. Four-year evaluation of the effect of vaccination against *Coxiella burnetii* on reduction of animal infection and environmental contamination in a naturally infected dairy sheep flock. *Appl Environ Microbiol.* (2011) 77:7405–7. doi: 10.1128/AEM.05530-11
21. Kersh GJ, Fitzpatrick KA, Self JS, Priestley RA, Kelly AJ, Lash RR, et al. Presence and persistence of *Coxiella burnetii* in the environments of goat farms associated with a Q fever outbreak. *Appl Environ Microbiol.* (2013) 79:1697–703. doi: 10.1128/AEM.03472-12
22. Angelakis E, Raoult D. Q fever. *Vet Microbiol.* (2010) 140:297–309. doi: 10.1016/j.vetmic.2009.07.016
23. Eldin C, Melenotte C, Mediannikov O, Ghigo E, Million M, Edouard S, et al. From Q fever to *Coxiella burnetii* infection: a paradigm change. *Clin Microbiol Rev.* (2017) 30:115–90. doi: 10.1128/CMR.00045-16
24. Melenotte C, Caputo A, Bechah Y, Lepidi H, Terras J, Kowalczywska M, et al. The hypervirulent *Coxiella burnetii* Guiana strain compared in silico, in vitro and in vivo to the Nine Mile and the German strain. *Clin Microbiol Infect.* (2019) 25:1155. doi: 10.1016/j.cmi.2018.12.039
25. Alonso E, López-Etxaniz I, Hurtado A, Liendo P, Urbaneja F, Aspritzaga I, et al. Q Fever outbreak among workers at a waste-sorting plant. *PLoS ONE.* (2015) 10:e0138817. doi: 10.1371/journal.pone.0138817
26. Alonso E, Eizaguirre D, López-Etxaniz I, Olaizola JI, Ocabo B, Barandika JF, et al. A Q fever outbreak associated to courier transport of pets. *PLoS ONE.* (2019) 14:e0225605. doi: 10.1371/journal.pone.0225605
27. Hurtado A, Alonso E, Aspritzaga I, López E I, Ocabo B, Barandika JF, et al. Environmental sampling coupled with real-time PCR and genotyping to investigate the source of a Q fever outbreak in a work setting. *Epidemiol Infect.* (2017) 145:1834–42. doi: 10.1017/S0950268817000796
28. Schets FM, de Heer L, de Roda Husman AM. *Coxiella burnetii* in sewage water at sewage water treatment plants in a Q fever epidemic area. *Int J Hyg Environ Health.* (2013) 216:698–702. doi: 10.1016/j.ijheh.2012.12.010
29. Huijsmans CJ, Schellekens JJ, Wever PC, Toman R, Savelkoul PH, Janse I, et al. Single-nucleotide-polymorphism genotyping of a *Coxiella burnetii* during a Q fever outbreak in The Netherlands. *Appl Environ Microbiol.* (2011) 77:2051–7. doi: 10.1128/AEM.02293-10
30. Glazunova O, Roux V, Freylikman O, Sekeyova Z, Fournous G, Tyczka J, et al. *Coxiella burnetii* genotyping. *Emerg Infect Dis.* (2005) 11:1211–7. doi: 10.3201/eid1108.041354

31. Barandika JF, Álvarez-Alonso R, Jado I, Hurtado A, García-Pérez AL. Viable *Coxiella burnetii* in hard cheeses made with unpasteurized milk. *Int J Food Microbiol.* (2019) 303:42–5. doi: 10.1016/j.ijfoodmicro.2019.05.010
32. Therneau T, Atkinson B. *rpart*: Recursive Partitioning and Regression Trees. R package version 4.1-15 (2019).
33. Milborrow S. *rpart.plot*: Plot 'rpart' Models: An Enhanced Version of 'plot.rpart'. R package version 3.0.8 (2019)
34. R Core Team. *A Language and Environment for Statistical Computing.* (2019).
35. Anderson AD, Szymanski TJ, Emery MP, Kohrs PH, Bjork AC, Marsden-Haug N, et al. Epizootiological investigation of a Q fever outbreak and implications for future control strategies. *J Am Vet Med Assoc.* (2015) 247:1379–86. doi: 10.2460/javma.247.12.1379
36. Pandit P, Hoch T, Ezanno P, Beaudeau F, Vergu E. Spread of *Coxiella burnetii* between dairy cattle herds in an enzootic region: modelling contributions of airborne transmission and trade. *Vet Res.* (2016) 47:48. doi: 10.1186/s13567-016-0330-4
37. Astobiza I, Barral M, Ruiz-Fons F, Barandika JF, Gerrikagoitia X, Hurtado A, et al. Molecular investigation of the occurrence of *Coxiella burnetii* in wildlife and ticks in an endemic area. *Vet Microbiol.* (2011) 147:190–4. doi: 10.1016/j.vetmic.2010.05.046
38. González-Barrio D, Ruiz-Fons F. *Coxiella burnetii* in wild mammals: a systematic review. *Transbound Emerg Dis.* (2019) 66:662–671. doi: 10.1111/tbed.13085
39. Berri M, Crochet D, Santiago S, Rodolakis A. Spread of *Coxiella burnetii* infection in a flock of sheep after an episode of Q fever. *Vet Rec.* (2005) 157:737–40. doi: 10.1136/vr.157.23.737
40. Reichel R, Mearns R, Brunton L, Jones R, Horigan M, Vipond R, et al. Description of a *Coxiella burnetii* abortion outbreak in a dairy goat herd, and associated serology, PCR and genotyping results. *Res Vet Sci.* (2012) 93:1217–24. doi: 10.1016/j.rvsc.2012.04.007
41. Rousset E, Berri M, Durand B, Dufour P, Prigent M, Delcroix T, et al. *Coxiella burnetii* shedding routes and antibody response after outbreaks of Q fever-induced abortion in dairy goat herds. *Appl Environ Microbiol.* (2009) 75:428–33. doi: 10.1128/AEM.00690-08
42. Joulié A, Rousset E, Gasqui P, Lepetitcolin E, Leblond A, Sidi-Boumedine K, et al. *Coxiella burnetii* circulation in a naturally infected flock of sheep: individual follow-up of antibodies in serum and milk. *Appl Environ Microbiol.* (2017) 83:e00222–17. doi: 10.1128/AEM.00222-17
43. Ruiz-Fons F, Astobiza I, Barandika JF, Juste RA, Hurtado A, García-Pérez AL. Measuring antibody levels in bulk-tank milk as an epidemiological tool to search for the status of *Coxiella burnetii* in dairy sheep. *Epidemiol Infect.* (2011) 139:1631–6. doi: 10.1017/S0950268810003134

44. Astobiza I, Tilburg JJ, Piñero A, Hurtado A, García-Pérez AL, Nabuurs-Franssen MH, et al. Genotyping of *Coxiella burnetii* from domestic ruminants in northern Spain. *BMC Vet Res*. (2012) 8:241. doi: 10.1186/1746-6148-8-241
45. Dal Pozzo F, Renaville B, Martinelle L, Renaville R, Thys C, Smeets F, et al. Single nucleotide polymorphism genotyping and distribution of *Coxiella burnetii* strains from field samples in Belgium. *Appl Environ Microbiol*. (2016) 82:81–6. doi: 10.1128/AEM.02799-15
46. Piñero A, Barandika JF, García-Pérez AL, Hurtado A. Genetic diversity and variation over time of *Coxiella burnetii* genotypes in dairy cattle and the farm environment. *Infect Genet Evol*. (2015) 31:231–5. doi: 10.1016/j.meegid.2015.02.006
47. de Bruin A, van der Plaats RQ, de Heer L, Paauwe R, Schimmer B, Vellema P, et al. Detection of *Coxiella burnetii* DNA on small-ruminant farms during a Q fever outbreak in the Netherlands. *Appl Environ Microbiol*. (2012) 78:1652–7. doi: 10.1128/AEM.07323-11
48. de Bruin A, Janse I, Koning M, de Heer L, van der Plaats RQ, van Leuken JP, et al. Detection of *Coxiella burnetii* DNA in the environment during and after a large Q fever epidemic in the Netherlands. *J Appl Microbiol*. (2013) 114:1395–404. doi: 10.1111/jam.12163
49. Carrié P, Barry S, Rousset E, de CR, Sala C, Calavas D, et al. Swab cloths as a tool for revealing environmental contamination by Q fever in ruminant farms. *Transbound Emerg Dis*. (2019) 66:1202–9. doi: 10.1111/tbed.13137