

BIBLIOGRAFIA:

1. Leblanc SJ (2020). Review: Relationships between metabolism and neutrophil function in dairy cows in the peripartum period. *Animal* 14(S1):44-54.
2. Leblanc (2004). Peripartum serum vitamin E, retinol, and beta-carotene in dairy cattle and their associations with disease: *J. Dairy Sci.* 87:609-619.
3. Duffield (2009). Impact of hyperketonemia in early lactation dairy cows on health and production : *J. Dairy Sci.* 92:571-580.
4. Dohoo (1984). Subclinical ketosis prevalence and associations with production and disease: *Can. J. Comp. Med.* 48: 1-5.
5. Ospina, P. A., D. V. Nydam, T. Stokol, and T. R. Overton (2010). Associations of elevated nonesterified fatty acids and beta-hydroxybutyrate concentrations with early lactation reproductive performance and milk production in transition dairy cattle in the Northeastern United States. *J. Dairy Sci.* 93:1596-1603.
6. Walsh R.B., Walton J. S. y col. (2007). The effect of subclinical ketosis in early lactation on reproductive performance of postpartum dairy cows *J. Dairy Sci*; 90: 2788-2796.
7. Leblanc (2010). Monitoring metabolic health of dairy cattle in the transition period. *J. Repro. Dev.* 56:S29-S35.
8. McArt et al. (2015). Hyperketonemia in early lactation dairy cattle: A deterministic estimate of component and total cost per case. *J. Dairy Sci.* 98 :2043-2054

