



Bibliografía

Referencias

- Arias, M., Lomba, C., Dacal, V., Vázquez, L., Pedreira, J., ..., Paz-Silva, A. (2011). Prevalence of mixed trematode infections in an abattoir receiving cattle from northern Portugal and north-west Spain. *Veterinary Record*, 168, 408. doi: 10.1136/vr.d85
- Calvo, J. (2021). Influencia del manejo en la prevalencia e intensidad de eliminación de los parásitos digestivos y pulmonares en rebaños de pequeños rumiantes en Galicia. Tesis Doctoral. Universidade de Santiago de Compostela.
- Cienfuegos, S., Díaz, P., Vázquez, L., Dacal, V., Lago, N., ..., López, C. (2009a). Prevalencia e intensidad de parasitación en granjas de pequeños rumiantes en Galicia. XIII Jornadas sobre producción animal (ITEA). Zaragoza: 143-145.
- Díaz, P., Lomba, C., Pedreira, J., Arias, M., Sánchez-Andrade, ..., Paz-Silva, A. (2006). Analysis of the IgG antibody response against paramphistomidae trematoda in naturally infected cattle. *Veterinary Parasitology*, 140, 281-288. doi:10.1016/j.vetpar.2006.04.007
- Díaz, P., Pedreira, J., Sánchez-Andrade, R., Suárez, J., Arias, M., ..., Paz-Silva, A. (2007). Risk periods of infection by *Calicophoron daubneyi* (digenea:Paramphistomidae) in cattle from oceanic climate areas. *Parasitology Research*, 101, 339-342. doi:10.1007/s00436-007-0493-z
- González-Warleta, M., Lladosa, S., Castro-Hermida, J. A., Martínez-Ibeas, A. M., Conesa, D., ..., Mezo, M. (2013). Bovine paramphistomosis in Galicia (Spain): Prevalence, intensity, aetiology and geospatial distribution of the infection. *Veterinary Parasitology*, 191, 252-263. doi:10.1016/j.vetpar.2012.09.006
- Mage, C., Bourgne, H., Toullieu, J.M., Rondelaud, D., Dreyfuss G. (2002). *Fasciola hepatica* and *Paramphistomum daubneyi*: changes in prevalences of natural infections in cattle and in *Lymnaea truncatula* from central France over the past 12 years. *Veterinary Research* 33, 439-447. Doi: 10.1051/vetres:2002030
- Mason, C., Stevenson, H., Cox, A., Dick, I. (2012). Disease associated with immature paramphistome infection in sheep. *Veterinary Record*, 170, 343-344. doi:10.1136/vr.e2368
- Millar, M., Colloff, A., Scholes, S. (2012). Bovine health: Disease associated with immature paramphistome infection. *Veterinary Record*, 171, 509-510. doi:10.1136/vr.e7738
- Orjales, I., Mezo, M., Miranda, M., González-Warleta, M., Rey-Crespo, F., ..., López-Alonso, M. (2017). Helminth infections on organic dairy farms in Spain. *Veterinary Parasitology*, 243, 115-118. doi:10.1016/j.vetpar.2017.06.031.
- Sanchís, J., Sánchez-Andrade, R., Macchi, M. I., Piñeiro, P., Suárez, J. L., ..., Arias, M. S. (2013). Infection by paramphistomidae trematodes in cattle from two agricultural regions in NW Uruguay and NW Spain. *Veterinary Parasitology*, 191, 165-171. doi:10.1016/j.vetpar.2012.07.028
- Toolan, D. P., Mitchell, G., Searle, K., Sheehan, M., Skuce, P. J., Zadoks, R. N. (2015). Bovine and ovine rumen fluke in Ireland—Prevalence, risk factors and species identity based on passive veterinary surveillance and abattoir findings. *Veterinary Parasitology*, 212, 168-174. doi:10.1016/j.vetpar.2015.07.040
- Vázquez, L., Dacal, V., Cienfuegos, S., Díaz, P., Lago, N., ..., López, C. (2008). Occurrence of trematode infections in sheep managed in a semiextensive system in northwestern Spain. *The XVI*

