

BIBLIOGRAFIA

Aguirre DH, Gaido AB, Cafrune MM, Castelli ME, Mangold AJ, Guglielmon AA (2005). Eprinomectin pour-on for control of *Boophilus microplus* (Canestrini) ticks (Acari: Ixodidae) on cattle. *Vet Parasitol* 127(2):157-63. doi: 10.1016/j.vetpar.2004.09.027.

Agencia Española de Medicamentos y Productos Sanitarios AEMPS (2023). CIMA.vet. Medicamentos Veterinarios. <https://cimavet.aemps.es/cimavet/publico/home.html>

Davey RB, George JE (2002). Efficacy of macrocyclic lactone endectocides against *Boophilus microplus* (Acari: Ixodidae) infested cattle using different pour-on application treatment regimes. *J Med Entomol* 39(5):763-9. doi: 10.1603/0022-2585-39.5.763.

del Castillo C, Pinedo R, Rodríguez L, Chávez A (2016). Evaluación de tres formulaciones comerciales de aplicación pour on bajo condiciones de campo y su efecto in vitro en el control de *Boophilus microplus* (Acari: Ixodidae) en Bovinos de Ceja de Selva. *Rev Inv Vet Perú* 27(1): 145-157. doi.org/10.15381/rivep.v27i1.11446.

do Nascimento CG, Bragaglia G, Toma SB, de Souza Magalhães V, Cid YP, Scott FB (2020). Injectable eprinomectin for cattle: Tick efficacy and pharmacokinetics. *J Vet Pharmacol Ther.* 43(2):171-178. doi: 10.1111/jvp.12840.

Ghoke SS, Bhikane AU, Awaz KB (2013). Therapeutic efficacy of eprinomectin pour-on against tick infestation - A report of 24 bovines. *Intas Polivet Vol. 14* (I): 72-73.

Lifschitz A, Nava S, Mangold A, Imperiale F, Ballent M, Canevari J, Lanusse C (2016). Eprinomectin accumulation in *Rhipicephalus* (*Boophilus*) *microplus*: Pharmacokinetic and efficacy assessment. *Vet Parasitol* 215: 11-16. doi.org/10.1016/j.vetpar.2015.11.005.

Lifschitz A, Nava S, Guglielmon AA, Imperiale F, Farias C, Mangold AJ, Lanusse C. (2008). Failure of ivermectin and eprinomectin to control *Amblyomma parvum* in goats: characterization of acaricidal activity and drug pharmacokinetic disposition. *Vet Parasitol* 156(3-4):284-92. doi: 10.1016/j.vetpar.2008.05.014.

Lohmeyer KH, Miller JA, Pound JM, Oehler DD (2009). Efficacy of eprinomectin and doramectin against *Amblyomma americanum* (Acari: Ixodidae) on cattle. *J Econ Entomol* 102(2):809-14. doi: 10.1603/029.102.0244.

Maciel WG, Lopes WDZ, Gomes LVC, Cruz BC, Felippelli G, Santos IBD, Borges FA, Gonçalves WA Junior, Scarpa AB, Nicaretta JE, Bastos TSA, da Costa AJ (2016). Susceptibility of *Rhipicephalus* (*Boophilus*) *microplus* to fluzaron (2.5mg/kg) and a combination of novaluron (2.0mg/kg) + eprinomectin (0.36mg/kg) in field studies in Brazil. *Prev Vet Med.* 135:74-86. doi: 10.1016/j.pvetmed.2016.10.019.

Nava S, Guglielmon AA (2009). Difficulties to control natural infestation with *Otobius megnini* (Acari: Argasidae) nymphs in cattle with systemic biocides. *Res Vet Sci.* 87(2):258-9. doi: 10.1016/j.rvsc.2009.02.012.

Nazir T, Katoch R, Godara R, Yadav A (2013). Efficacy of eprinomectin pour-on against *Rhipicephalus* (*Boophilus*) *microplus* on buffaloes. *J Parasit Dis* 37(2):166-7. doi: 10.1007/s12639-012-0152-x.

Rodríguez-Vivas RI, Arieta-Román RJ, Pérez-Cogollo LC, Rosado-Aguilar JA, Ramírez-Cruz GT, Basto-Estrella, G (2010). Uso de lactonas macrocíclicas para el control de la garrapata *Rhipicephalus* (*Boophilus*) *microplus* en el ganado bovino. *Arch Med Vet* 42, 115-123.

Strydom T, Lavan RP, Torres S, Heaney K (2023). The economic impact of parasitism from nematodes, trematodes and ticks on beef cattle production. *Animals* (Basel) 13(10):1599. doi: 10.3390/ani13101599.

