

BIBLIOGRAFIA

Cañas C., R., Quiroz, R. A., León-Velarde, C., Posadas, A., & Osorio, J. (2003). Quantifying energy dissipation by grazing animals in harsh environments. *Journal of Theoretical Biology*, 225(3), 351–359. [https://doi.org/10.1016/S0022-5193\(03\)00260-1](https://doi.org/10.1016/S0022-5193(03)00260-1)

Fernández, C., Hernández, A., Gomis-Tena, J., & Llor, J. J. (2021). Changes in nutrient balance, methane emissions, physiologic biomarkers, and production performance in goats fed different forage-to-concentrate ratios during lactation. *Journal of Animal Science*, 99(7), 1–13. <https://doi.org/10.1093/jas/skab114>

Lachica, M., & Aguilera, J. F. (2003). Estimation of energy needs in the free-ranging goat with particular reference to the assessment of its energy expenditure by the 13C-bicarbonate method. *Small Ruminant Research*, 49(3), 303–318. [https://doi.org/10.1016/S0921-4488\(03\)00146-9](https://doi.org/10.1016/S0921-4488(03)00146-9)

Lachica, M., & Aguilera, J. F. (2005). Energy needs of the free-ranging goat. *Small Ruminant Research*, 60(1-2 SPEC. ISS.), 111–125. <https://doi.org/10.1016/j.smallrumres.2005.06.006>

Lachica, M., Somlo, R., Barroso, F. G., Boza, J., & Prieto, C. (1999). Goats locomotion energy expenditure under range grazing conditions: Seasonal variation. *Journal of Range Management*, 52(5), 431–435. <https://doi.org/10.2307/4003768>

Marutsova, V., & Binev, R. (2017). Body condition score, nonesterified fatty acids and beta-hydroxybutyrate concentrations in goats with subclinical ketosis. *Agricultural Science and Technology*, 9(4), 282–285. <https://doi.org/10.15547/ast.2017.04.053>

NRC, National Research Council, USA. (1981). *Nutrient Requirements of Goats: Angora, Dairy, and Meat Goats in Temperate and Tropical Countries*. Washington, D.C.: Ed.National Academy Press.

Osuji, P. O. (1974). The Physiology of Eating and the Energy Expenditure of the Ruminant at Pasture. *Journal of Range Management*, 27(6), 437. <https://doi.org/10.2307/3896717>

Ríos, C., Marín, M. P., Catafau, M., & Wittwer, F. (2006). Concentraciones sanguíneas de β -hidroxibutirato, NEFA, colesterol y urea en cabras lecheras de tres rebaños con sistemas intensivos de producción y su relación con el balance nutricional. *Archivos de Medicina Veterinaria*, 38(1), 19–23. <https://doi.org/10.4067/s0301-732x2006000100003>

Singh, M., & Ludri, R. S. (2002). Milk production, blood metabolites and circulatory levels of hormones in crossbred goats. *Asian-Australasian Journal of Animal Sciences*, 15(7), 963–967. <https://doi.org/10.5713/ajas.2002.963>

Zamuner, F., DiGiacomo, K., Cameron, A. W. N., & Leury, B. J. (2020). Short communication: Associations between nonesterified fatty acids, β -hydroxybutyrate, and glucose in periparturient dairy goats. *Journal of Dairy Science*, 103(7), 6672–6678. <https://doi.org/10.3168/jds.2019-17163>

